

Spatial Distribution of Medusa *Cunina octonaria* and Frequency of Parasitic Association with *Liriope tetraphylla* (Cnidaria: Hydrozoa: Trachylina) in Temperate SW Atlantic Waters

Francisco Alejandro Puente-Tapia^{1,*}, Florencia Castiglioni², Gabriela Failla Siquier², and Gabriel Genzano¹

¹*Instituto de Investigaciones Marinas y Costeras (IIMyC), Facultad de Ciencias Exactas y Naturales, Universidad Nacional de Mar del Plata (FCEyN, UNMdP), Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Mar del Plata, Argentina.*

**Correspondence: E-mail: alecspuente@gmail.com (Puente-Tapia). Tel +5492236938742. E-mail: genzanogabriel@gmail.com (Genzano)*

²*Laboratorio de Zoología de Invertebrados, Departamento de Biología Animal, Facultad de Ciencias, Universidad de la República, Montevideo, Uruguay. E-mail: f.castiglioni17@gmail.com (Castiglioni); gabrielafailla@gmail.com (Failla Siquier)*

Received 16 April 2020 / Accepted 27 August 2020 / Published xx October 2020

Communicated by Ruiji Machida

This study examined the spatial distribution of the medusae phase of *Cunina octonaria* (Narcomedusae) in temperate Southwestern Atlantic waters using a total of 3,288 zooplankton lots collected along the Uruguayan and Argentine waters (34°–56°S), which were placed in the Medusae collection of the Universidad Nacional de Mar del Plata, Argentina. In addition, we reported the peculiar parasitic association between two hydrozoan species: the polypoid phase (stolon and medusoid buds) of *C. octonaria* (parasite) and the free-swimming medusa of *Liriope tetraphylla* (Limnomedusae) (host) during one-year sampling period (February 2014 to March 2015) on the coasts of Mar del Plata, Argentina. We examined the seasonality, prevalence, and intensity of parasitic infection. Metadata associated with the medusa collection was also used to map areas of seasonality where such association was observed. *Cunina octonaria* was found from southern Uruguay to coasts of Mar del Plata (34.8–38.2°S, 57.2–54.0°W), with the highest abundances and frequency of occurrence in the Río de la Plata estuary. The parasitic association was identified during the austral warm period (spring-summer season) and up to mid-autumn. Out of 21,734 *L. tetraphylla* specimens that were examined, 316 were parasitized (prevalence = 1.5%) exclusively in the manubrium and gastric peduncle, with an infection intensity of 1 to 2 stolons per host. Furthermore, the medusoid buds per stolon ranged from 11 and 29 at different stages of development. No significant differences were observed between the umbrella diameter of

parasitized and non-parasitized *L. tetraphylla* specimens, nor was any significant correlation identified between umbrella diameter and prevalence, and intensity of infection. According to the aggregation coefficient, *C. octonaria* had an overdispersed distribution in the host population. All parasitized hosts showed stomach vacuity due to the location of the stolon, which blocked the mouth of the host. We identified the parasitic association in coasts of Mar del Plata, as well as in both coasts of the Río de la Plata Estuary (Uruguayan-Argentinean coasts). In the Southwestern Atlantic, several biological interactions between medusae and other groups have been identified; however, the specific host selectivity of *C. octonaria* for *L. tetraphylla* was not previously identified. Here we discuss the ecological importance of this association during the holoplanktonic life history of the narcomedusae. Additionally, we report the southern limit of the spatial distribution of this particular parasitic association in the Southwestern Atlantic, thus increasing the knowledge of biological associations of gelatinous zooplankton (Cnidaria and Ctenophora) on Uruguayan and Argentinean coasts.

Key words: Argentine Sea, Stolon phase, Medusoid buds, Limnomedusae, Narcomedusae.

Citation: Puente-Tapia FA, Castiglioni F, Siquier GF, Genzano G. 2020. Spatial distribution of medusa *Cunina octonaria* and frequency of parasitic association with *Liriope tetraphylla* (Cnidaria: Hydrozoa: Trachylina) in temperate SW Atlantic waters. Zool Stud 59:0mm. doi:-.

BACKGROUND

Medusae (*i.e.*, Medusozoa, which includes Hydro-, Scypho-, Cubo-, and Stauromedusae) (see Marques and Collins 2004; Lewis Ames 2018) are an abundant group distributed in a wide spectrum of aquatic ecosystems, including marine-coastal environments (Larson 1986; Suchman and Sullivan 2000) and some continental freshwaters systems (Arbačiauskas and Lesutienė 2005), occurring in all latitudes (Mills 1995), from polar to tropical waters, whereas in the water column they can be found from the surface to abyssal depths (Vinogradov 1968), including benthic and planktonic environments (Miranda et al. 2016). One of the ecological roles of this group includes a great variety of biological interactions with several groups such as protists, cnidarians, trematodes, mollusks, crustaceans, echinoderms, fish, among others (Ohtsuka et al. 2009 and references therein).

The study of the medusae community from temperate Southwestern Atlantic waters (SWAw) (southern Brazil, Uruguay, and Argentina) has allowed us to know the hydromedusae faunal composition, the spatial distribution (mainly of the most abundant and frequent species)

(Genzano et al. 2008; Oliveira et al. 2016; Dutto et al. 2019), and some of the above mentioned interactions (e.g., Sal Moyano et al. 2012; Díaz Briz et al. 2012 2017). However, knowledge of the spatial distribution of some species, such as *Cunina octonaria* McCrady, 1959 (Trachylina: Narcomedusae: Cuninidae) is scarce. The present study represents the first attempt to describe its spatial distribution along this vast area of the SWAw, as well as the association of this species and *Liriope tetraphylla* (Chamisso and Eysenhardt, 1821) (Trachylina: Limnomedusae: Geryoniidae), which has not been previously reported for SWAw. We should highlight this particular parasitic relationship, due to two species of hydromedusae are involved, each one in a specific stage of its life cycle: the free-swimming medusa phase of *L. tetraphylla* as host to the polypoid phase (stolon and medusa buds) of *C. octonaria* as parasite. In addition, although the polypoid phase and the free-swimming medusa of *C. octonaria* are the same species, each one has different reproductive strategies and energy requirements.

Hydrozoans are a group of cnidarians with complex and diverse life cycles, as well as reproductive strategies that result in a wide range of diversity of polyps, colonies, and medusa morphologies, as well as complete loss or reduction of the polyp or medusa stage in some species (Cartwright and Nawrocki 2010). Morphological and molecular data have allowed us to classify to Hydrozoa into two sister groups: Hydroidolina and Trachylina (Bouillon et al. 2006). Trachylina includes the most enigmatic cnidarians, as they differ from other hydrozoans in their morphology, development, and life cycles (Collins et al. 2008). This group includes four orders: Limnomedusae, Narcomedusae, Trachymedusae, and Actinulida. Order Limnomedusae is a small group with dimorphic benthic-pelagic cycle, characterized by a small or even nonexistent polyp in some species (Bouillon et al. 2004). When there is no polyp, its embryo directly transforms into the medusa (e.g., *L. tetraphylla*) (Osadchenko and Kraus 2018). For Narcomedusae species, the development is indirect and do not include a real intermediate sessile polyp stage (Mayer 1910), their connection to organic, benthic substrates is lost and perform their entire life cycle in the pelagic realm (Bouillon 1987; Bentlage et al. 2018). However, there is another way to establish an association with substrates: by using another living organism during a larval stage (Bouillon 1987), which may develop in a sessile stage that lives in parasitic association with other cnidarians (e.g., *C. octonaria*) (Bouillon 1987; Osborn 2000).

This type of associations may be necessary for some species at a certain phase of their life-cycles and may, or may not, present host specificity. The larval stages of several Narcomedusae (such as *C. octonaria*) parasitize the gastrovascular system and manubrium of other medusae, using their host's food and tissues (Boero and Bouillon 2005; Raskoff and Robison 2005). Once grown, they leave their hosts and live freely (Boero and Bouillon 2005).

Although medusae have a wide range of morphologies and complex life history patterns (Bouillon and Boero 2000), few published studies describe the association between hydromedusae and other cnidarians. The study of biological associations has implications for marine biodiversity (Purcell and Arai 2001). Forming an association with medusae can enhance the population or diversity of the commensals since medusae provide shelter, act as food (Masuda et al. 2008), and protect the larval stages from predators (Ohtsuka et al. 2009). Analyzing this intragroup association (*i.e.*, intrahydrozoan), allows to see aspects of its population dynamic, life history, life cycle variation (Benovic et al. 1987; Giagrande et al. 1994; Lucas and Reed 2009), and ecological role (*e.g.*, trophic and ecological niches) (Marcogliese 2005). These aspects are essential to understand how populations develop and persist through the year (Graham et al. 2001; Lucas and Reed 2009). Parasites may be useful indicators of food-web structure and function because they depend on the presence of other hosts in their life cycles for transmission (Marcogliese 2005). There are numerous advantages of using parasites as indicators of trophic processes and food structure; for example, identifying direct trophic links between the host and other organisms, host trophic level, and predators (Marcogliese 2005).

MATERIALS AND METHODS

Spatial distribution of *Cunina octonaria*

To analyze the historical distribution of *C. octonaria* in temperate SWAw, we examined 3,288 zooplankton samples of the Medusae collection (MedCol) of the J.J. Nágera Coastal Station, Universidad Nacional de Mar del Plata (UNMDP), Argentina were revised. The collection includes zooplankton samples taken over 36 years (from 1980 to 2016) along the Uruguayan and Argentine continental shelves (34–56°S, 52–70°W), comprising an area ~7 million km² (Fig. 1). The sampling was done using fishery research vessels operated by the Instituto Nacional de Investigaciones y Desarrollo Pesquero (INIDEP, Argentina), as well as local research programs designed by the Instituto de Investigaciones Marinas y Costeras (IIMyC-CONICET, Mar del Plata, Argentina) and the Instituto Argentino de Oceanografía (IADO-CONICET/UNS, Bahía Blanca, Argentina). A variety of plankton nets (Bongo, Nackthai, Motoda, Pairovet, Calvet, and Multinet) were used in oblique trawls from the proximity of the bottom to the surface. The medusae were sorted, counted, identified, and the abundances were calculated and expressed as ind.m³ of filtered waters. The occurrence of *C. octonaria* was plotted based on its presence/absence in the set of samples. The spatial distribution of *L. tetraphylla* has been previously described in Dutto et al. (2019).




Fig. 1. Map of the study area: temperate Southwestern Atlantic waters. Yellow circle indicate the area studied for seasonality variation (Mar del Plata Harbor, Argentina, 38°08'17"S-57°31'18"W); red crosses represent the zooplankton samples of the dataset from the Medusae collection. RPE (Rio

de la Plata estuary); BsAs (Buenos Aires); SB (Samborombón Bay); SCT (San Clemente del Tuyú); MdP (Mar del Plata); BB (Bahía Blanca); SMG (San Matías Gulf); VP (Valdés Peninsula); SJG (San Jorge Gulf); TdF (Tierra del Fuego); M/F (Malvinas/Falkland Islands). The fine lines represent the 50, 100, 200, and 1000 m isobaths of the region. Scale bar = 250 km.

To analyze and compare the possible differences between austral warm (spring-summer) and cold periods (autumn-winter) regarding the frequency of occurrence and abundances of *C. octonaria* along with the geographical distribution, the data set was divided into two periods: from October to March (spring-summer) and from April to September (autumn-winter). Then, we plotted the presence/absence data and abundances of both periods; we performed Pearson's chi-square test to test the possible differences between both periods and Student *t*-test for the abundances differences (Sokal and Rohlf 1999).

Parasitic association

During the analysis of the temporal variation of the GZ species on the Mar del Plata Harbor (Argentina) (38°08'17"S–57°31'18"W) (Fig. 1), we identified the parasitic association between the polypoid phase of *C. octonaria* and the free-swimming medusa of *L. tetraphylla*; therefore, we used these sampling to described the seasonality of this association. The zooplankton sampling was obtained during a one-year sampling period (February 2014 to March 2015). The samples were collected with a weekly or biweekly frequency during the warm period and a monthly frequency in the cold-period with oblique tows using a standard zooplankton net (mouth diameter: 75 cm; mesh size: 500- μ m) and a flowmeter attached to the net mouth, which allowed us to calculate the volume of filtered water to estimate the numerical abundance (ind.m³) of the GZ. We first analyzed this parasitic association by describing the temporal variation of the abundances of the involved species. For *L. tetraphylla*, we reviewed the previous reports of Puente Tapia and Genzano (2019), and for *C. octonaria* we carried out several analyses: a one-way ANOVA to test if abundances of the medusa phase were significantly different between seasons. If significant differences were present, a *post hoc* Tukey's HSD test was performed to examine which seasons showed differences (Zar 1999). Data were previously normalized by logarithmic transformation to fulfill the assumption to the homogeneity of variances (Levene test). The analyzed specimens were stored at the MedCol with the tag *Cunina-Liriope-001*.

The total number of parasitized organisms per sample was recorded, as well as the number of stolons on each host specimens. With these data, the (a) parasitic prevalence and (b) intensity of infection were described. We determined these infection levels first in terms of the total values (*i.e.*,

considering the period of occurrence of the association) and later for each month. According to Bush et al. (1997): (a) prevalence is the number of hosts infected with one or more individuals of polypoid structure of *C. octonaria* per sample, divided by the number of hosts and expressed as a percentage; (b) intensity of infection is the number of stolons of *C. octonaria* in a single infected host. In addition, we also recorded the number of medusoid buds of *C. octonaria* in each stolon.

The total umbrellar diameter (UD, in mm) of the hosts was measured under a stereomicroscope to determine the relationship between host size and parasite load (considering only the number of stolons). Thus, the specimens were immersed in water in order to adopting a similar position that is observed in marine environment. A general size-frequency histogram per host was constructed to analyze the occurrence of the different host sizes with the presence/absence of polypoid phase of *C. octonaria*. The length classes were arbitrarily defined by applying the Sturges' rule (Zar 1999). Possible differences between the sizes of parasitized and non-parasitized hosts were analyzed by a Student-test (Sokal and Rohlf 1999).

After checking for the normality of data distribution and homogeneity of variance, we used Pearson's correlation coefficient (r) (Zar 1999) to analyze the relationship between the host size and prevalence and intensity of infection for each size group. After calculating the coefficient (r) and evaluating its significance, a Student's t -distribution test was applied. Considering the number of medusoid buds per host and using the r -coefficient, we examined the correlation of each size group. The level of statistical significance was $P \leq 0.05$ for all analyses. The assumption of normality of data (Kolmogorov-Smirnov test) and homogeneity of variance (Levene test) were verified by using the previous normalization with logarithmic transformation of the data (Zar 1999). To estimate the distribution of *C. octonaria* in the host population, we calculated the aggregation coefficient (k) of the negative binomial distribution. This analysis determined whether the host species had a normal, random or overdispersed distribution (Morales and Pino 1987).

The concept of host range (Rohde 2005) was used to classify parasite species as specialists or generalists. According to this concept, a specialist parasite species has a marked affinity for a specific host family, genus or species, while a generalist is one that parasitizes hosts from a number of different taxa. For this analysis, we reviewed the available literature on the associations between genus *Cunina* and other hosts species; We examined the gut content of the all *L. tetraphylla* organisms (including parasitized and non-parasitized) in terms of vacuity (*i.e.*, presence or lack of stomach content), which could be an indicator of the negative effects *C. octonaria* caused *L. tetraphylla*. Subsequently, the metadata associated with the MedCol served to determine: a) the areas of the spatial co-occurrence of *L. tetraphylla* and *C. octonaria* and b) geographical areas and the seasonality where this parasitic association was observed.

RESULTS

Spatial distribution of *Cunina octonaria* in temperate Southwestern Atlantic waters

According to the data of the MedCol, this narcomedusa was found in 28 of 3,288 zooplankton samples (0.85%). It had a spatial distribution from 34.8° to 38.2°S and 57.5° to 54.0°W (i.e., from southern Uruguayan coasts to waters of central-northern Buenos Aires Province, Argentina) (Fig. 2A). The main occurrence was observed in the estuarine area of the Río de la Plata. All samples showing *C. octonaria* specimens were collected at a depth lower than 50 m. Although, 92.85% of these samples ($n = 26$) were collected in the warm period and only 7.14% ($n = 2$) in the cold period, no significant differences were observed ($\chi^2 = 2.0$, $d.f. = 1$, $P = 0.16$) regarding the frequency of occurrence between both periods. In the cold period, the specimens were collected only in San Clemente del Tuyú (~36°S–56.8°W) and Mar del Plata coasts (~38°S–57.5°W), with abundances that ranged from 0.04 to 1.08 ind.m³ (Fig. 3A). In the warm period, the abundances ranged between 0.003 and 28.05 ind.m³, showing the highest abundances ~130 km south of the Samborombón Bay, followed by the Río de la Plata estuarine area (close to Uruguay) (Fig. 3B). No significant differences were found between the abundances from the warm and cold periods ($t = 0.89$, $d.f. = 24$, $P = 0.38$). *Liriope tetraphylla* showed a southern limit up to ~38°50'–40°S, ~60–61°W (~Bahía Blanca estuary, the adjacent shelf of El Rincón and Monte Hermoso, Buenos Aires Province, Argentina) (Fig. 2B) (see Dutto et al. 2019).




Fig. 2. Spatial distribution in temperate Southwestern Atlantic of the free-swimming medusa phase of *Cunina octonaria* (A) and *Liriope tetraphylla* (B) (including parasitized and non-parasitized specimens). Orange cycles represent the samples of zooplankton with species of medusa; black crosses represent the lack of individuals; (C) Areas of the association between the polypoid phase of *C. octonaria* and *L. tetraphylla* (red cycles). Acronyms in figure 1. Scale bars = 250 km.




Fig. 3. Comparison of the spatial distribution of the abundances (ind.m³) of the free-swimming medusa phase of *Cunina octonaria* between the (A) austral cold (autumn-winter) and (B) warm periods (spring-summer) in temperate Southwestern Atlantic waters. Red arrows (2A) indicate the regions where in which abundances were recorded during the cold period. Acronyms in figure 1. Scale bars = 250 km.

Analysis of parasitic association

In samples collected around the year (February 2014 to March 2015) in Mar del Plata Harbor, the free-swimming medusa of *C. octonaria* was observed only in the spring and summer seasons, with abundances ranging from 0.003 (October) to 3.82 ind.m³ (February) (Table 1). In terms of seasonality, no significant differences were observed in the abundance values (ANOVA: $F = 1.62$; $P = 0.34$). *Cunina octonaria* was found in salinities ranging from 33.7 to 36, while the water temperature oscillated between 14° and 22.3°C. *Liriope tetraphylla* occurred year-round, with the highest abundances during the warm period and the lowest in the cold seasons. Their monthly abundances ranged from 0.01 ind.m³ (May) to 80.98 ind.m³ (March) (Table 1). Significant differences were observed between climate seasons (ANOVA: $F = 14.30$, $P = 0.01$), particularly between autumn-spring (Tukey's HSD test = -1.70, $P = 0.01$), autumn-summer (Tukey = -2.28, $P < 0.01$), and winter-summer (Tukey's HSD test = -1.50, $P = 0.01$), *i.e.*, between the cold and warm periods.

Table 1. Abundance values (ind.m³) of the free-swimming medusa phase of *Cunina octonaria* and *Liriope tetraphylla* during the study year (February 2014 to March 2015) in Mar del Plata Harbor, Argentina

	<i>Cunina octonaria</i>	<i>Liriope tetraphylla</i>	Water temperature (°C)	Salinity
2014				
February	0.29	10.90	22.0	35.0
March	0.08	80.98	19.1	35.0
April	0.04	0.27	15.3	35.0
May	0	0.01	14.2	35.0
June	0	0.08	11.7	34.0
July	0	0.41	10.4	35.0
August	0	1.45	9.8	35.0
September	0	0.26	11.8	34.0
October	0.003	4.22	14.0	33.7
November	0	0.73	16.3	36.0
December	1.16	19.53	18.8	34.0
2015				
January	0.73	20.99	19.3	35.3
February	3.82	5.22	21.2	36.0
March	0	19.95	22.3	36.0

A total of 18 gelatinous species were identified (14 hydromedusae, 1 scyphomedusae, and 3 ctenophores), out of which the polypoid phase of *C. octonaria* was only observed in association with *L. tetraphylla*. The parasitic association was found during the spring and summer seasons, with water temperatures ranging from 14.0 to 22.0°C, and salinity from 34.0 to 36.0. Out of 21,734 individual of *L. tetraphylla* analyzed during the period of co-occurrence of the association, 316 were infected (total prevalence = 1.45%). All hosts had a single stolon structure (intensity of infection) with medusoid buds, except for one that had two stolons. The number of medusoid buds per host ranged from 11 (December) to 29 (January). These values varied monthly (Table 2).

Although, the umbrellar diameter (UD) of *L. tetraphylla* ranged from 0.5 to 14.3 mm, parasitized specimens oscillated from 1.7 to 7.0 mm. Medusae between 5 and 6 mm of the size-frequency histogram showed the highest number of parasitized specimens, with prevalence values of 2.5 and 3.5%, respectively (Fig. 4). No significant differences were observed between the UD of parasitized and non-parasitized specimens ($t = 1.0$, $d.f. = 281$, $P = 0.30$). No significant correlations between the UD and prevalence were observed ($r = 0.02$, $n = 244$, $P = 0.72$), nor between UD and intensity of infection (number of stolons). However, a significantly positive correlation between the number of medusoid buds per host and the UD was observed ($r = 0.61$, $n = 73$, $P < 0.01$, $t = 0.16$, $d.f. = 71$, $P < 0.05$): *i.e.*, the number of medusoid buds increased as the UD of the *L. tetraphylla* increased.

Table 2. Period of co-occurrence and parasitological parameters (prevalence and intensity of infection) of the association between *Liriope tetraphylla* (*Lt*) and *Cunina octonaria* (*Cu*) at Mar del Plata Harbor, including measurements of physicochemical and biological parameters

Period of co-occurrence	Total number of <i>Lt</i> analyzed	Total number of <i>Lt</i> parasitized	Prevalence (%)	Intensity of infection	Number of medusoid bud of <i>Cu</i> Min-Max	Surface water temperature (°C)	Salinity
February 2014	336	7	2.1	1	11-14	22.0	35.0
March 2014	14,757	44	0.3	1	12-21	19.1	35.0
October 2014	1925	2	0.1	1	11-13	14.0	33.7
December 2014	549	1	0.2	1	12-18	18.8	34.0
January 2015	4,167	262	6.3	1-2	13-39	21.2	36.0
Total	21,734	316	1.5	2			

The host species showed a parasite overdispersed distribution according to the aggregation coefficient ($k = 0.32$). That is, most of the examined organisms harbored few or no parasites. In contrast, few host individuals contained a large number of this narcomedusa.

Observation *in vivo* showed that the stolon of *C. octonaria* had several medusoid buds in different growth stages, which protruded from the characteristic manubrium and gastric peduncle of *L. tetraphylla*. A section of the stolon was inside the manubrium and another one on the outside (Fig. 5A-D). All stolon structures had medusoid buds. The stolon had an elongated and cylindrical shape widening in the basal region to the interior of the manubrium, while the external side had flat or oval shapes (Fig. 5C-D). Newly liberated medusae and free-swimming medusa of *C. octonaria* are shown in figure 5E-F.

In addition to the Mar del Plata, we identified the association in San Clemente del Tuyú (Argentina) (36°21'S-56°43'W), but Castiglioni and Faila Siquier (personal observations) also identified it in Maldonado, Uruguay (34°38'S-54°5'W). These finding indicate that the association is present on both coast of the Río de la Plata estuarine area (Uruguay-Argentina) (Fig. 2C). On Mar del Plata coasts, the association was observed in the austral spring (October and December) and summer (from January to March), while in San Clemente del Tuyú, it was observed during the mid-autumn (mid-April and mid-May). Finally, on the Uruguayan coasts, it was identified from January to March (summer to early autumn).

A general analysis of the gut content of all individuals of *L. tetraphylla* during the occurrence of the association showed that the stomachs of the parasitized individuals were completely empty, whereas some of the non-parasitized organisms had chaetognaths, crustaceans,

and detritus. This vacuity might be due to the location of the polypoid phase, which prevented food intake.




Fig. 4. Percentage of parasitized and non-parasitized *Liriopse tetraphylla* for the different size classes. Grey bars: pooled data of non-parasitized hosts; black bars: specimens parasitized with polypoid phase of *Cunina octonaria*.




Fig. 5. Parasitic association between the polypoid phase of *Cunina octonaria* (1) (parasite) and *Liriope tetraphylla* (host) from Mar del Plata, Argentina. (A-D) Location of the stolon in the manubrium and gastric peduncle of *L. tetraphylla*; (D) detail of the stolon and the medusoid buds; (E) newly liberated medusae from the stolon; (F) comparison between newly liberated medusa (2) and adult individual (3) of *C. octonaria*. Lowercase letters indicate the structure of the host species: m (manubrium); mo (mouth); su (subumbrella); t (tentacles); eu (exumbrella). Scale bars: A–C and F = 1 mm; D–E = 500 μ m.

DISCUSSION

Spatial distribution of *Cunina octonaria* and *Liriope tetraphylla* in temperate Southwestern Atlantic waters

Cunina octonaria is a Narcomedusae widely distributed in the Pacific, Atlantic, and Indian Oceans (Burke 1975; Segura-Puertas 1984; Galea 2007; Segura-Puertas et al. 2009; Andrade Ruíz 2010; Oliveira et al. 2016; Schuchert 2020). In the SWAw it has been identified from Colombia to Argentina (Oliveira et al. 2016). In the present study, we described the distribution in temperate waters, going from the southern Uruguayan to Mar del Plata coasts ($\sim 38^{\circ}$ S) as the southern limit, with the higher abundances and frequency of occurrence in the Río de la Plata estuary and adjacent

waters. However, according to Zamponi (1983) isolated records have shown its presence near of the Malvinas/Falkland Islands ($\sim 51^{\circ}46'S-59^{\circ}31'W$). Although we reviewed more than 3,200 zooplankton samples, from which $\sim 20\%$ ($n = 647$) were distributed under this latitude, *C. octonaria* was not found in the samples from the region indicated by Zamponi (1983); therefore, his finding could not be confirmed.

During the study year in Mar del Plata, *C. octonaria* was found in temperatures between 14° and $22.3^{\circ}C$. According to Soares Moreira (1978), in laboratory conditions *C. octonaria* shows little tolerance to low temperatures, with an ideal value of $15^{\circ}C$. At $10^{\circ}C$ the specimens crumpled, while at $5^{\circ}C$ they became completely quiet. Therefore, it is possible that the Patagonian and Sub-Antarctic regions are unfavorable zone for *C. octonaria* due to the low temperatures, mainly in the austral cold period.

During the cold period, the shelf circulation of the temperate SWA consists of the northward flow cold waters of the Malvinas/Falkland Current, which is characterized by low-salinity and nutrient-rich waters. In contrast, during the warm period, the influence is from the southward tropical-subtropical flow warm waters of the Brazil Current, characterized by oligotrophic and salty waters (Piola and Matano 2001; Piola et al. 2010). According to Vannucci (1957) and Navas-Pereira (1973), *C. octonaria* is a typical species of the Brazil Current. Earlier records confirm this to be a widely distributed frequent species in warm waters (Kramp 1965); consequently, the fact that this narcomedusa was only found in the warm period and, sporadically at the early autumn, could be due passively transportation through the warm current. In this context, the contribution of the Río de la Plata waters are a natural barrier to the distribution of different marine organisms (Boltovskoy et al. 1999). However, this barrier is intermittent because in periods of low flow and with the effects of northeastern winds, it is possible to find species, which usually live in northern areas of Argentina (Uruguay and Brazil), in southern regions and localities such as Mar del Plata (Mianzan et al. 2001; Mianzan and Acha 2008).

On the other hand, several studies have focused on *L. tetraphylla* because it is the most frequent taxon on temperate SWAw (Dutto et al. 2019). This species was mainly found in the Río de la Plata estuary and surroundings areas (Fig. 4B). During the study year in Mar del Plata, *L. tetraphylla* was observed in temperatures between 9.8° and $22.3^{\circ}C$ and salinity ranging from 33.7 to 36 (Puente Tapia and Genzano 2019). According to Vannucci (1957), this species showed an ideal values of temperature from $20^{\circ}C$ to $25^{\circ}C$ and 35-36 of salinity. However, it was observed in strongly diluted surface waters at the mouth of the Amazon river. The spatial distribution of *L. tetraphylla* was observed up to \sim Bahía Blanca estuary, the adjacent shelf of El Rincón and Monte Hermoso coasts (Buenos Aires Province, Argentina). Historically, this medusa has been identified as a dominant species in the zooplankton community in some productive areas of the Argentine Sea

(Zamponi and Genzano 1994; Sorarrain 1998); it reaches high abundances up to 4.7×10^6 ind.m³ (Mianzan et al. 2000). Despite that *L. tetraphylla* is one of the most abundant and common medusa in temperate SWAw (Mianzan and Guerrero 2000; Mianzan et al. 2000; Gaitán 2004; Failla Siquier 2006; Dutto et al. 2017), the parasitic association with *C. octonaria* had not been previously reported in the region.

Host selectivity

Analyzing the parasitic association over a year in Mar del Plata allowed us to observe a host specificity of *L. tetraphylla*. We took into account some factors to try to identify the reason behind this selectivity in this South Atlantic region: A) the parasite feeding, B) spatial distribution, C) abundances, and D) swimming behavior of the host, as well as E) the characteristic manubrium and gastric peduncle of *L. tetraphylla*. These factors are discussed below; however, these aspects arose when we compared our observations with the background information.

A) *Parasite feeding strategy*: host specificity is often the result of eco-biological factors such as the parasite feeding (Rohde and Rohde 2005). We found feeding similarities between the polypoid phase of *C. octonaria* and its medusa phase; thus, analyzing both phases might explain the selectivity. Some parasites infect many hosts, but only those which have similar feeding habits (Rohde and Rohde 2005). Although *L. tetraphylla* feeds on crustaceans, chaetognaths (Puente Tapia personal observations), fish eggs, and larvae (Mianzan et al. 2012), Narcomedusae seem to feed primarily on soft-bodied prey such as pelagic individuals (Purcell and Mills 1988). However, in *Cunina*, the parasitic polypoid phase not only absorbs the reproductive tissues of the host (gelatinous individual) but also ingests food captured by the host (Raskoff and Robison 2005). Therefore, this polypoid structure share a similar diet with the medusa phase, feeding on gelatinous organisms, with an additional food source to “steal” from the host. According to Lima and Valentin (2001), the associations may be that of a simple guest (only feed on the food taken by the host) or as a parasitism if the parasite uses the tissue of the host as a source of nutrition. The polypoid of *C. octonaria* exhibits both feeding strategies.

B) *Spatial distribution of the host*: The spatial distribution of the parasites may be determined by the host distribution (Marcogliese 2002). Both *C. octonaria* and *L. tetraphylla* share almost the same spatial distribution, with a southern limit of co-occurrence on the Mar del Plata coasts. In Argentina Sea, some of the host species previously identified for *C. octonaria* have been reported (Table 3), which shown a southernmost limit distribution (Patagonian and Sub-Antarctic regions) with respect to *L. tetraphylla* (Rodríguez 2012). Despite that, *C. octonaria* was absent on these coasts, suggesting a close relationship with *L. tetraphylla*. There is no “universal” parasite that

infects all available microhabitats on all available host species. In other words, each parasite species occupies a particular niche: it is microhabitat- and host-specific (Marcogliese 2005).

Although *C. octonaria* is considered an oceanic species (Johnson and Allen 2005), in the temperate SWA it was located at a depth lower than 50 m, that is, shallow waters. Although some *L. tetraphylla* individuals were observed between the 50 m and 100 m isobaths, a high proportion of the positive samples were located at a depth lower than 20 m (Fig. 2B). Certain ecological patterns emerge from the distribution of parasites in marine waters. Horizontal gradients exist in parasite diversity, with more species in shallow waters as a result of the wide range of hosts available (Marcogliese 2002). In pelagic waters, parasite species richness declines with depth and then increases in the bottom waters (Marcogliese 2005 and references therein).

Table 3. List of species of the genus *Cunina* in parasitic association with other hydromedusae as hosts

Parasite species	Host species			Reference
	Order	Family	Species	
<i>Cunina octonaria</i>	Anthomedusae	Cytaeidae	<i>Cytaeis tetrastyla</i>	9,10
		Protiaridae	<i>Halitiara inflexa</i>	9,10
		Teissieridae	<i>Teissiera medusifera</i>	10
		Oceaniidae	<i>Turritopsis nutricula</i>	2,5,6,13
	Leptomedusae	Laodiceidae	<i>Laodicea undulata</i>	7,10
		Campanulariidae	<i>Clytia hemisphaerica</i>	11
	Trachymedusae	Rhopalonematidae	<i>Aglaura hemistoma</i>	13
			<i>Aglaura</i> sp.	2
			<i>Rhopalonema velatum</i>	13
	Limnomedusae	Geryoniidae	<i>Liriope tetraphylla</i>	2,9,10,12,13
<i>Geryonia proboscidalis</i>			3,7,10,13	
Narcomedusae	Cuninidae	<i>Cunina octonaria</i>	1,13	
<i>Cunina peregrina</i>	Trachymedusae	Rhopalonematidae	<i>Aglaura hemistoma</i>	7,10,13
			<i>Rhopalonema velatum</i>	3,7,10,13
	Narcomedusae	Cuninidae	<i>Cunina octonaria</i>	8
			<i>Cunina peregrina</i>	13
			Solmarisidae	<i>Pegantha martagon</i>
	Solmundaeginidae	<i>Solmundella bitentaculata</i>	10	
<i>Cunina proboscidea</i>	Limnomedusae	Geryoniidae	<i>Geryonia proboscidalis</i>	4,13
<i>Cunina becki</i>	Narcomedusae	Cuninidae	<i>Cunina peregrina</i>	10,13

1) Mayer (1910); 2) Vannucci (1957); 3) Kramp (1959); 4) Kramp (1961); 5) Calder (1968); 6) Calder (1971); 7) Bouillon (1978); 8) Goy (1979); 9) Bouillon et al. (1986); 10) Bouillon (1987); 11) Zhenzu and Huiduan (1994); 12) present study; 13) references cited in Bouillon (1987).

C) *Host abundances*: *C. octonaria* and *L. tetraphylla* showed a co-occurrence during the warm period. During this period, *L. tetraphylla* had high abundances (Mianzan et al. 2000), which is a factor that could facilitate the encounter between the two species (in this instance, medusa

phase of *L. tetraphylla* and larval stage of *C. octonaria*) and begin the parasitic association. Being extremely diffuse, the marine environment forms unique barriers that theoretically could prevent parasite infection. In some cases, parasites are typically short-lived (Marcogliese 2005) and needs to rapidly find the host individual. The highest abundances of *L. tetraphylla* occur in the warm period and coincide spatiotemporally with *C. octonaria*, which might represent an advantage for the parasite phase of this narcomedusa and help it penetrate the host and continue its life cycle. However, during the warm period of the study year, we identified other medusae with similar or even higher abundances than *L. tetraphylla*, such as the leptomedusae *Obelia* sp. and *Eucheilota ventricularis* McCrady, 1859 on the coasts of Argentina (Puente-Tapia and Genzano 2019), and *Eucheilota maculata* Hartlaub, 1894 and *Clytia hemisphaerica* (Linnaeus, 1767) on the coasts of Uruguay (Failla Siquier personal observation). A question arose from observing these medusae: Why was *L. tetraphylla* the only host species? The ability to infect a wide range of hosts further serves parasites to survive in an otherwise unforgiving habitat (Marcogliese 2005).

D) *Swimming behavior of the host*: Parasitism on hydromedusae is not unusual, and the slow movement of many species provides pelagic or benthic parasites to take residence for either locomotion or protection (Lucas and Reed 2009). According to Osborn (2000), the swimming behavior of the medusae might favor contact with parasites and increase the encounter rate. The feeding strategy of *L. tetraphylla* involves spreading its tentacles and remaining still in that position wait for its preys (Mills 1981; Puente Tapia personal observations). This posture could make it easier for the larvae of *C. octonaria* to be pulled or sucked into the stomach cavity or gastric system (including the manubrium), where they may remain during their development.

In general, we observed two types of feeding-swimming strategies of the most abundant species during the study year: filter-feeding (*Obelia* sp. and *E. ventricularis*) and ambush predation (*L. tetraphylla*) (Sutherland et al. 2016). In the filter-feeding, the species has a continuous pulse of the umbrella, creating water currents towards their mouth (Boero et al. 2007), which could prevent *C. octonaria* from entering into the umbrella cavity. In ambush predator, the species remains in apparent quiet for at least some seconds while waiting for their prey (Puente Tapia personal observations).

Some parasite stages may be transmitted usually swim and search out a susceptible host and penetrate it. Those forms that are passively transmitted are usually ingested by the appropriate host in the life cycle for transmission to occur (Marcogliese 2005). In case of medusae, the ciliated planula larva has been interpreted as a motile stage (Bouillon and Boero 2000), which lives only from a few hours to a few days (Bouillon 1994). Therefore, it is essential to analyze the possible chemical signals of the host species to determine if the larva is attracted. Thereon, infective stages exhibit further morphological and behavioral adaptations that increase the probability of parasitism.

Those that are passively transmitted may resemble typical prey of the target host. Actively transmitted stages display behaviors such as phototropism, geotropism, or timing of hatch or emergence that serve to place them in the target host habitat (Williams and Jones 1994). These stages may be chemically attracted to the host over small spatial scales (Marcogliese 2005).

E) *Manubrium and gastric peduncle of the host*: The peculiar manubrium and the cylindrical gastric peduncle of *L. tetraphylla*, which is longer than umbrella diameters (Bouillon et al. 2004) are possibly some of the most visible morphological differences between *L. tetraphylla* and the other medusae species identified in the study area. For example, the leptomedusae *Obelia* sp. and *E. ventricularis* have a small manubrium or lack gastric peduncle (*Obelia* sp.) (Bouillon et al. 2004). But, Is the manubrium one of the determining factors of host selectivity in the region? The manubrium and gastric peduncle of the previously identified host species of *C. octonaria* display various forms and sizes: *Turritopsis nutricula* McCrady, 1857 has a manubrium large; *Aglaura hemistoma* Péron and Lesueur, 1810 has a small manubrium and a somewhat shorter gastric peduncle than subumbrella radius; *Rhopalonema velatum* Gegenbaur, 1857 has a manubrium narrow reaching extended almost velar opening; *Geryonia proboscidalis* (Forsskål, 1775) is almost similar to *L. tetraphylla* (Bouillon et al. 2004); while *C. octonaria* lack a manubrium (Bouillon et al. 2004, 2006). Therefore, it seems that the manubrium is not the determining factor of the host selectivity.

Other remarks

The parasitic association was observed in three different areas: Maldonado (Uruguay), San Clemente del Tutú and Mar del Plata coasts (Argentina). However, in different sectors of the Brazilian coasts, the larval phase of *Cunina* sp. parasitizes *T. nutricula* and *Proboscidactyla ornata* (McCrady, 1859); two larvae of *C. octonaria* were observed on *Aglaura* sp. (Vannucci 1957); while in the São Sebastião Channel (Brazil; Tropical SWAw), the manubrium of *L. tetraphylla* was observed with larvae of *Cunina* (Migotto 2020).

After the reviewing the available literature, we found four different species of the genus *Cunina* in parasitic association with other medusae. For *C. octonaria* have been identified 11 species. Hence, this review allowed us classify *C. octonaria* as a generalist species according to the concept of host range (Rohde 2005) (Table 3). Other three species of the genus *Cunina* have been found to parasite medusae: *C. peregrina* Bigelow, 1909 was observed in association with six species, while for *C. proboscidea* Metschnikoff and Metschnikoff, 1871 and *C. becki* Bouillon, 1985 have exhibit only one host species each until now. Although parasites are often generalists, they are linked to particular types of host functional groups or feeding guilds. Members of such

guilds share lifestyles, diet preferences, and depth ranges. Consequently, members of these guilds often share a similar parasite fauna whose constituent species follow common transmission pathways (Marcogliese 2002).

Moreover, in the present study a single stolon of *C. octonaria* per host (only one specimen had two stolons), but Bouillon (1987) described specimens parasitized by up to three different stolons in a single host (two was the most common number). We could not analyze all negative effects of the polypoid of *C. octonaria* on *L. tetraphylla*; however, we observed that all parasitized organisms had an empty stomach, possibly due to the location of the stolon in the mouth opening, which prevented *L. tetraphylla* from feeding.

The distribution of *C. octonaria* in *L. tetraphylla* populations agrees with the aggregate distribution pattern that is typical in parasite species (Bush et al. 2001). Most parasite species are not randomly or uniformly distributed in host populations; they show an aggregated (overdispersed) distribution, that is, some individuals of host populations are more heavily infected than expected in a random distribution, and others are lightly infected or not at all (Rohde 1984). Several factors contribute to the aggregation distribution: a single parasite individual may multiply on or in the host, as observed in the *C. octonaria* life cycle. This distribution pattern stabilizes the host-parasite association. Some biological function of this parasitic aggregation could be considered as an effect “in favor” of the host population. Since few hosts were heavily infected, only they could die. The death of these hosts could ensure the completion of the life cycle of a parasite that depends on being eaten by the next host, but the host population as a whole is not greatly affected (Rohde 1984 and references therein).

CONCLUSIONS

The life cycle with indirect metagenetic and parasitic development of *Cunina octonaria* (Bouillon 1987), has some advantages, for example, the use as substrate a free swimming medusa, allows it to have access to additional food sources (Raskoff and Robison 2005), as well as greater spatial dispersion compared to a species with benthic polyp (Boero et al. 2008). The diversity and plasticity of the “traditional” life forms and life history modes across Medusozoa means that medusae can colonize a plethora of ecosystems (Lewis Ames 2018 and references therein). Based on previous observations it has been concluded that the advantages to parasitism include the brooding, protecting and enhancing the survival of young by providing a safe, nutritive, and internal habitat for development (Osborn 2000).

An important factor that limits the number of hosts could be the availability of suitable host species (Rohde and Rohde 2005). The Uruguayan and Argentine waters are characterized by high species diversity, including multiple corporal forms, life cycles, seasonality, abundances, feeding strategies, and swimming behavior. However, taking into account the parasite feeding, spatial distribution, abundances, swimming behavior of *L. tetraphylla*, and its characteristic manubrium and gastric peduncle, it is difficult to explain the host selectivity toward *L. tetraphylla* in this temperate region. The combination of two or more of mentioned factors could explain the host selectivity of *C. octonaria*. Therefore, extensive research using molecular markers or chemical analysis is needed to prove this specificity (Ohtsuka et al. 2009).

Temperate SWAw represent a favorable region both *C. octonaria* and *L. tetraphylla* to complete their particular life cycles. The food availability in this region meets the energy requirements of both species. The medusa phase of *C. octonaria* could benefit from the large amount of available food, to eat and reproduce, while their larva stage could use the high abundances of *L. tetraphylla* to facilitate the encounter, start the parasitic phase and the subsequent feeding and reproductive advantages. Thus, a chain of hydro-eco-biological events is observed, in which each species benefits from some of these factors to complete their life cycle.

Our dataset indicates that the Argentine Sea represent the southern limit of the distribution of this interaction in the SWAw. Finally, this study increases the knowledge about biological associations between GZ and other groups living on the Uruguayan and Argentinean coasts. However, it is necessary to conduct more zooplankton studies to identify the exact moment when the interaction begins, possible chemical-molecular signals between host and larva of *C. octonaria*, the effects on the host, and other regions where the association could exist.

Acknowledgments: We thank the collaboration of the Oceanographer C. Brelles and the anonymous reviewers for their comments, which greatly improved this manuscript. The authors thanks to R. Ascención and Biol. AP. Vázquez-Vázquez for their help with the English grammar and style of this manuscript. This work was supported by the UNMdP, Argentina (grant number EXA 829/17).

Authors' contributions: FAPT and GG designed the study, performed the field work, and wrote the manuscript. FAPT performed the data analysis. FC and GFS provided the biological data from Uruguayan coasts, contributed to the analysis of data and to organize the manuscript. All authors participated in the discussion of the results, revising the manuscript, and approved the final version.

Competing interests: The authors declare that they have no conflicts of interest.

Availability of data and materials: The materials used in the present study are available for anybody's research in the future. Specimens used here are deposited in the MedCol of the UNMDP, Argentina.

Consent for publication: Nor applicable.

Ethic approval consent to participate: Nor applicable.

REFERENCES

- Andrade Ruíz EC. 2010. Sistemática y ecología de las medusas (Cnidarias: Hydrozoa) en la zona costera sur de la Bahía de Santa Elena durante el periodo (octubre 2004-octubre 2005). Master dissertation, Universidad de Guayaquil.
- Arbačiauskas K, Lesutienė J. 2005. The freshwater jellyfish (*Craspedacusta sowerbii*) in Lithuanian waters. *Acta Zool Litu* **15**:54–57. doi:10.1080/13921657.10512609.
- Benovic A, Dubravko J, Bender A. 1987. Enigmatic changes in the hydromedusan fauna of the northern Adriatic Sea. *Nature* **326**:597–600. doi:10.1038/326597a0.
- Bentlage B, Osborn KJ, Lindsay DJ, Hopcroft RR, Raskoff KA, Collins AG. 2018. Loss of metagenesis and evolution of a parasitic life style in a group open ocean jellyfish. *Mol Phylogenetics Evol* **124**:505–559. doi:10.1016/j.ympev.2018.02.030.
- Boero F, Bouillon J. 2005. Cnidaria and Ctenophora (cnidarians and comb jellies). *In*: Rohde K (ed) *Marine parasitology*, 1st edn. CSIRO Publishing and CABI, Melbourne and Wallingford.
- Boero F, Bucci C, Colucci AMR, Gravili C, Stabili L. 2007. *Obelia* (Cnidaria, Hydrozoa, Campanulariidae): a microphagous, filter-feeding medusa. *Mar Ecol* **28**:178–183. doi:10.1111/j.1439-0485.2007.00164.x
- Boero F, Bouillon J, Gravili C, Miglietta MP, Parsons T, Piraino S. 2008. Gelatinous plankton: irregularities rule the world (sometimes). *Mar Ecol Prog Ser* **356**:299–310. doi:10.3354/meps07368.
- Boltovskoy D, Gibbons MJ, Hutchings L, Biner D. 1999. General biological features of the South Atlantic. *In*: Boltovskoy D (ed) *South Atlantic Zooplankton*. Backhuys Publishers, Leiden.
- Bouillon J. 1978. Hydroméduses de l'Archipel des Séchelles et du Mozambique. *Rev Zool Adr* **92**:117–172.

- Bouillon J. 1987. Considérations sur le développement des Narcoméduses et sur leur position phylogénétique. *Indo-Malayan Zool* **4**:189–278.
- Bouillon J. 1994. Embranchement des Cnidaires (Cnidaria). In: Grassé PP, Doumenc D (eds) *Traité de Zoologie, Cnidaires et Cténares* Tome III, Fascicule 2. Masson, Paris, France.
- Bouillon J, Claereboudt M, Seghers G. 1986. Hydroméduses de la baie de Hansa (Mer de Bismarck; Papouasie Nouvelle-Guinée). Répartition, conditions climatiques et hydrologiques. *Indo-Malayan Zool* **3**:105–152
- Bouillon J, Boero F. 2000. Phylogeny and classification of hydroidomedusae. *The Hydrozoa: a new classification in the light of old knowledge. Thal Salent* **24**:1–296
- Bouillon J, Medel MD, Pagès F, Gili JM, Boero F, Gravili C. 2004. Fauna of the Mediterranean Hydrozoa. *Sci Mar* **68**:5–438.
- Bouillon J, Gravili C, Pagès F, Gili JM, Boero F. 2006. An introduction to Hydrozoa. *Mém Mus Natl Hist Nat* **194**:1–591.
- Burke DW. 1975. Pelagic Cnidaria of Mississippi Sound and adjacent waters. *Gulf Res Rep* **5**:23–38.
- Bush AO, Lafferty KD, Lotz JM, Shostak AW. 1997. Parasitology meets ecology on its own terms: Margolis et al. revisited. *J Parasitol* **83**:575–583. doi:10.2307/3284227.
- Bush AO, Fernández JC, Esch GW, Seed Jr. 2001. *Parasitism: The Diversity and Ecology of Animal Parasites*. Cambridge University Press, Cambridge, UK.
- Calder DR. 1968. Hydrozoa of the Southern Chesapeake Bay. PhD dissertation, Virginia Institute of Marine Science.
- Calder DR. 1971. Hydroids and Hydromedusae of southern Chesapeake Bay. *Vir Inst Mar Sci Spec Paper* **1**:1–125.
- Cartwright P, Nawrocki AM. 2010. Character Evolution in Hydrozoa (Phylum Cnidaria). *Integr Comp Biol* **50**:456–472. doi:10.1093/icb/icq089.
- Collins AG, Bentlage B, Lindner A, Lindsay D, Haddock SHD, Jarms G, Norenburg JL, Jankowski T, Cartwright P. 2008. Phylogenetics of Trachylina (Cnidaria: Hydrozoa) with new insights on the evolution of some problematical taxa. *J Mar Biol Assoc UK* **88**:1673–1685. doi:10.1017/S0025315408001732.
- Díaz-Briz LM, Martorelli SR, Genzano GN, Mianzan HW. 2012. Parasitism (Trematoda, Digenea) on medusae from the southwestern Atlantic Ocean: medusa hosts, parasite prevalences, and ecological implications. *Hydrobiologia* **690**:215–226. doi:10.1007/s10750-012-1040-1.
- Díaz-Briz L, Sánchez F, Marí N, Mianzan H, Genzano G. 2017. Gelatinous zooplankton (ctenophores, salps and medusae): an important food resource of fishes in the temperate SW Atlantic Ocean. *Mar Biol Res* **13**:630–644. doi:10.1080/17451000.2016.1274403.

- Dutto MS, Genzano GN, Schiariti A, Lecanda J, Hoffmeyer MS, Pratolongo P. 2017. Medusae and ctenophores from Bahía Blanca Estuary and neighboring inner shelf (Southwest Atlantic Ocean, Argentina). *Mar Biodivers Rec* **10**:1–14. doi:10.1186/s41200-017-0114-1.
- Dutto MS, Chazarreta CJ, Rodríguez CS, Schiariti A, Diaz-Briz LM, Genzano GN. 2019. Macroscale abundance patterns of hydromedusae in the temperate Southwestern Atlantic (27°–56°S). *PLoS ONE* **14**:e0217628. doi:10.1371/journal.pone.0217628.
- Failla Siquier MG. 2006. Zooplankton gelatinoso de la costa uruguaya. In: Menafrá R, Rodríguez-Gallego L, Scarabino F, Conde D (eds) *Bases para la Conservación y el manejo de la costa uruguaya*. Vida Silvestre Uruguay, Montevideo.
- Galea HR. 2007. Hydroids and hydromedusae (Cnidaria: Hydrozoa) from the fjords region of southern Chile. *Zootaxa* **1597**:1–116. doi:10.11646/zootaxa.1597.1.1.
- Gaitán EN. 2004. Distribución, abundancia y estacionalidad de *Liriope tetraphylla* (Hidromedusae) en el Océano Atlántico Sudoccidental y su rol ecológico en el estuario del Río de la Plata. Dissertation, Universidad Nacional de Mar del Plata.
- Genzano G, Mianzan H, Bouillon J. 2008. Hydromedusae (Cnidaria: Hydrozoa) from the temperate southwestern Atlantic Ocean: a review. *Zootaxa* **1750**:1–18. doi:10.11646/zootaxa.1750.1.1.
- Giagrande A, Geraci S, Belmonte G. 1994. Life-cycle and life-history diversity in marine invertebrates and the implications in community dynamics. *Ocean Mar Biol* **32**:305–333.
- Goy J. 1979. Méduses. *Res Sci Camp de la Calypso* **11**:263–296.
- Graham WM, Pagès F, Hamner WM. 2001. A physical context for gelatinous zooplankton aggregations: a review. *Hydrobiologia* **451**:199–212. doi:10.1023/A:1011876004427.
- Johnson W, Allen DM. 2005. *Zooplankton of the Atlantic and Gulf coasts: a guide to their identification and ecology*. The Johns Hopkins University Press. Baltimore.
- Kramp PL. 1959. The Hydromedusae of the Atlantic Ocean and adjacent waters. *Dana Rep* **46**:1.283.
- Kramp PL. 1961. Synopsis of the medusae of the world. *J Mar Biol Assoc UK* **40**:1–469.
- Kramp PL. 1965. The Hydromedusae of the Pacific and Indian Ocean. *Dana Rep* **63**:1–161.
- Larson RJ. 1986. Water content, organic content and carbon and nitrogen composition of medusae from the northeast Pacific. *J Exp Mar Biol Ecol* **99**:107–120. doi:10.1016/0022-0981(86)90231-5.
- Lima MCG, Valentin JL. 2001. New records of Amphipoda Hyperiididae in associations with gelatinous zooplankton. *Hydrobiologia* **448**:229–235. doi:10.1023/A:1017593120143.
- Lewis Ames C. 2018. Medusa: A review of an ancient Cnidarian body form. In: Kloc M, Kubiak J. (eds) *Marine organisms as model systems in biology and medicine. Results and Problems in cell differentiation*, 1st edn. Springer, Cham.

- Lucas CH, Reed AJ. 2009. Observations on the life histories of the narcomedusae *Aeginura grimaldii*, *Cunina peregrina* and *Solmissus incisa* from the western North Atlantic. *Mar Biol* **156**:373–379. doi:10.1007/s00227-008-1089-6.
- Marcogliese DJ. 2002. Food webs and the transmission of parasites of marine fish. *Parasitology* **124**:S83–S99. doi:10.1017/s003118200200149x.
- Marcogliese DJ. 2005. Transmission of marine parasites. In: Rohde K (ed) *Marine parasitology*, 1st edn. CSIRO Publishing and CABI, Melbourne and Wallingford.
- Marques AC, Collins AG. 2004. Cladistic analysis of Medusozoa and cnidarian evolution. *Invertebr Biol* **123**:23–42.
- Masuda R, Yamashita Y, Matsuyama M. 2008. Jack mackerel *Trachurus japonicus* juveniles use jellyfish for predator avoidance and as a prey collector. *Fisheries Sci* **74**:276–284. doi:10.1111/j.1444-2906.2008.01522.x.
- Mayer AG. 1910. *Medusae of the world I-II*. Carnegie Institution of Washington, Washington.
- Mianzan HW, Acha E. 2008. Procesos ecológicos en el Mar Patagónico. In: Estado de Conservación del Mar Patagónico y Áreas de Influencia. Puerto Madryn. Available via <http://marpatagonico.org>. Accessed 30 June 2020.
- Mianzan HW, Guerrero RA. 2000. Environmental patterns and biomass distribution of gelatinous macrozooplankton. Three study cases in the South-western Atlantic Ocean. *Sci Mar* **64**:215–224. doi:10.3989/scimar.2000.64s1215.
- Mianzan HW, Sorarrain D, Burnett J, Lutz L. 2000. Mucocutaneous junctional and flexural parasthesias caused by the holoplanktonic Trachymedusae *Liriope tetraphylla*. *Dermatology* **201**:46–48. doi:10.1159/000018429.
- Mianzan HW, Collini R, Schiariti A, Suarez J, Acha EM. 2012. Impacto de la depredación por medusas (*Liriope tetraphylla* y *Eucheilota ventricularis*) sobre huevos de peces: experiencias de laboratorio. *INIDEP*:1-12.
- Mianzan HW, Lasta C, Acha EM, Guerrero R, Machi G, Bremec C. 2001. The Río de la Plata Estuary, Argentina-Uruguay. In: Seeliger U, Kjerve B (eds) *Coastal Marine Ecosystems of Latin America*. Ecological Studies (analysis and synthesis), 1st edn. Springer, Berlin.
- Miranda LS, Hirano YM, Mills CE, Falconer A, Fenwick D, Marques AC, Collins AG. 2016. Systematics of stalked jellyfishes (Cnidaria: Sturozoa). *PeerJ* **4**:e1951. doi:10.7717/peerj.1951.
- Migotto AEM. 2020. Hydromedusa. Cifonauta image database. Center for Marine Biology of University of São Paulo. <http://cifonauta.cebimar.usp.br/media/1615/>. Accessed 30 June 2020.
- Mills CE. 1981. Seasonal occurrence of planktonic medusae and ctenophores in the San Juan Archipelago (NE Pacific). *Wasmann J Biol* **39**:6–29.

- Mills CE. 1995. Medusae, siphonophores, and ctenophores as planktivorous predators in changing global ecosystems. *ICES J Mar Sci* **52**:575–581. doi:10.1016/1054-3139(95)80072-7.
- Morales G, Pino LA. 1987. *Parasitología Cuantitativa*. Fundación Fondo Editorial, Acta Científica Venezolana, Caracas.
- Navas-Pereira D. 1973. Análise qualitativa e quantitativa do zooplâncton. Realatório sobre a segunda pesquisa oceanográfica e pesqueira do Atlântico Sul entre Torres e Maldonado (Lat 29°-35°S). Programa Rio Grande do Sul II. GEDIP e Instituto Oceanográfico, Univ São Paulo **3**:29–92.
- Ohtsuka S, Koike K, Lindsay D, Nishikawa J, Miyake H, Kawahara M, Mulyadi, Mujiono N, Hiromi J, Komatsu H. 2009. Symbionts of marine medusae and ctenophores. *Plankton Benthos Res* **4**:1–13. doi:10.3800/pbr.4.1
- Oliveira OM, Miranda TP, Araujo EM, Ayón P, Cedeño-Poso CM, Cepeda-Mercado AA, Córdoba P, Cunha AF, Genzano GN, Haddad MA, Mianzan HW, Migoto AE, Miranda LS, Morandini AC, Nagata RM, Nascimento KB, Nogueira Jr. M, Palma S, Quiñones J, Rodriguez CS, Scarabino F, Schiariti A, Stampar SN, Tronolone VB, Marques AC. 2016. Census of Cnidaria (Medusozoa) and Ctenophora from South American marine waters. *Zootaxa* **4194**:1–256. doi:10.11646/zootaxa.4194.1.1.
- Osadchenko BV, Kraus YuA. 2018. Trachylina: The group that remains enigmatic despite 150 years of investigations. *Russ J Dev Biol* **49**:134-145. doi:10.1134/S1062360418030074.
- Osborn DA. 2000. Cnidarian “Parasites” on *Solmissus incisa*, a Narcomedusa. *Sci Mar* **64**:157–163. doi:10.3989/scimar.2000.64s1157.
- Piola AR, Matano RP. 2001. Brazil and Falkland (Malvinas) Currents. *In*: Steele JH, Thorpe SA, Turekian KK (eds) *Encyclopedia of Ocean Sciences*, 2nd edn. Academic Press, London.
- Piola AR, Palma ED, Piola AR. 2010. The influence of the Brazil and Malvinas Currents on the Southwestern Atlantic Shelf circulation. *Ocean Sci* **6**:983–995. doi:10.5194/os-6-983-2010.
- Puente-Tapia FA, Genzano G. 2019. Seasonal succession of gelatinous zooplankton (medusae and ctenophores) from Mar del Plata Harbor, Argentina (SW Atlantic Ocean). *Ecol Austral* **29**:339–351. doi:10.25260/EA.19.293.0.880.
- Purcell JE, Mills CE. 1988. The correlation between nematocyst types and diets in pelagic Hydrozoa. *In*: Hessinger D, Lenhoff H (eds) *The biology of nematocysts*, 1st edn. Academic Press, San Diego, USA.
- Purcell JE, Arai MN. 2001. Interaction of pelagic cnidarians and ctenophores with fish: a review. *Hydrobiologia* **451**:27–44. doi:10.1023/A:1011883905394.

- Raskoff KA, Robinson BH. 2005. A novel mutualistic relationship between a doliolid and a cnidarian, *Bythotiara dolioeques* sp. nov. J Mar Biol Assoc UK **85**:583–593. doi:10.1017/s0025315405011513.
- Rodríguez CS. 2012. Hidromedusas del Atlántico sudoccidental: Biodiversidad y patrones de distribución. PhD dissertation, Universidad Nacional de Mar del Plata.
- Rohde K. 1984. Ecology of marine parasites. Helgoländer Meeresunters **37**:5–33.
- Rohde K. 2005 (ed). Marine parasitology. CSIRO Publishing and CABI, Melbourne and Wallingford.
- Rohde K, Rohde PP. 2005. The ecological niches of parasites. In: Rohde K (ed) Marine parasitology, 1st edn. CSIRO Publishing and CABI, Melbourne and Wallingford.
- Sal Moyano MP, Schiariti A, Giberto DA, Díaz Briz LM, Gavio MA, Mianzan HW. 2012. The symbiotic relationship between *Lychnorhiza lucerna* (Scyphozoa, Rhizostomeae) and *Libinia spinosa* (Decapoda, Epialtidae) in the Río de la Plata (Argentina-Uruguay). Mar Biol **159**:1933–1941. doi:10.1007/s00227-012-1980-z.
- Segura Puertas L. 1984. Morfología, sistemática y zoogeografía de las medusas (Cnidaria: Hydrozoa y Scyphozoa) del Pacífico Tropical Oriental. Anales del Instituto de Ciencias del Mar y Limnología, UNAM **8**:1–320.
- Segura-Puertas L, Celis L, Chiaverano L. 2009. Medusozoans (Cnidaria: Cubozoa, Scyphozoa and Hydrozoa) of the Gulf of Mexico. In: Felder DL, Capm DK (eds) Gulf of Mexico, Origin, waters and biota. Vol. 1. Biodiversity, 1st edn. Texas A&M University Press.
- Schuchert P. 2020. World Hydrozoa Database. Accessed through: World Register of Marine Species (WoRMS). <http://www.marinespecies.org/aphia.php?p=taxdetails&id=22754>. Accessed 30 June 2020.
- Soares Moreira G. 1978. A preliminary laboratory study on the salinity and temperature tolerance of some medusae from the São Paulo coast, Brazil. Bolm Inst Oceanograr S Paulo **27**:45–55. doi:10.1590/S0373-55241978000200002.
- Sokal R, Rohlf F. 1999. Introducción a la Bioestadística. Editorial Reverté, S.A., Barcelona.
- Sorarrain D. 1998. Cambios estacionales en la biomasa de organismos gelatinosos en relación a otros zooplancteres en la Bahía Samborombón. Dissertation, Universidad Nacional de Mar del Plata.
- Suchman CL, Sullivan BK. 2000. Effect of prey size on vulnerability of copepods to predation by the scyphomedusae *Aurelia aurita* and *Cyanea* sp. J Plank Res **22**:2289–2306. doi:10.1093/plankt/22.12.2289.

- Sutherland KR, Gemmell BJ, Colin SP, Costello JH. 2016. Prey capture by the cosmopolitan hydromedusae, *Obelia* spp., in the viscous regime. *Limnol Oceanogr* **61**:2309–2317. doi:10.1002/lno.10390.
- Vannucci M. 1957. On Brazilian Hydromedusae and their distribution in relation to different water masses. *Bol Inst Ocean* **8**:23–109.
- Vinogradov FE. 1968. Vertical distribution of the Oceanic zooplankton. Nauka, Moscow.
- Williams H, Jones A. 1994. Parasitic Worms of Fish. Taylor and Francis, Bristol.
- Zamponi MO. 1983. Ecología de las hidromedusas del Mar Epicontinental argentino. *Neotrópica* **29**:65–81.
- Zamponi MO, Genzano G. 1994. Seasonal distribution of hydromedusae from Samborombón Bay (Buenos Aires, Argentina). *Plankton Newsletter* **19**:51–56.
- Zar JH. 1999. Biostatistical Analysis. Prentice-Hall, Upper Saddle River, New Jersey.
- Zhenzu X, Huiduan W. 1994. The life cycle of parasitic species of *Cunina octonaria* McCrady from Xiamen Harbor, China. *J Xiamen Univ Natur Sci* **33**:154–159.